

Web: www.AgilelV.com | Tel: (201) 751-2202 | Fax: (201) 266-0437 | Email: info@AgilelV.com

LEQEMBI MEDICATION ORDER

DOB
Patient's weight in pounds
\Box G30.1 Alzheimer's with late onset (at \geq 65y/o)
☐ G30.8 Other Alzheimer's disease
nent are required. (Usually it must be mild for approval.)
history, including relevant tests and laboratory results, all available UMS and MoCA), differential diagnoses (ie. DLB, FTD, etc.), and patientes.
twelve (12) months?
via PET or CSF. If CSF, document why was PET not obtained.
e provided prior to the $5^{ m th}, 7^{ m th}$ and 14 $^{ m th}$ infusions.
(ARIA). Testing for and clinical evaluation regarding ARIA before and during remains the <u>sole</u> responsibility of the ordering provider. The MRI reports provided before the start of each round of therapy.
ed, documentation of pt education re increased ARIA risk must be provided.)
onal therapy such as Aduhelm? ently under control?
include documentation of counseling that use of Leqembi with such therapy ent and/or guardian has shared in the decision-making to undergo Leqembi
National Patient Registry trial number:
tions to be administered 15 minutes prior to the start of the infusion treatment. nasonemg
ote: Only a single course can be selected per order form.
☐ 10mg/kg IV every 2 weeks for treatments number 14-20
☐ For treatments #21 and beyond only:
10mg/kg IV every 2 weeks for treatments
sed over 1 hour. The IV line shall have a 0.2 micron in-line filter attached. Post infusion toms at start, throughout infusion, and after completion.
Reaction ema, nausea, vomiting, abdominal pain, hypotension, and respiratory distress. at 50 ml/hr. Call ordering provider to report reaction. ne, methylprednisolone, albuterol and oxygen as needed. call 911. Repeat if severe symptoms persist.
Name: Date:
Indiv. NPI #: License:
I de troma de la companya de la comp

_____ Direct Phone to Contact Person: ___

Best Contact Person in Office: _